

Calderdale and Huddersfield NHS Foundation Trust

West Yorkshire and Harrogate A19

The Calderdale and Huddersfield NHS Foundation Trust incorporates Calderdale Royal Hospital in Halifax and Huddersfield Royal Infirmary http://www.cht.nhs.uk/>http://www.cht.nhs.uk/>http://www.cht.nhs.uk/>http://www.cht.nhs.uk/>

Patient inclusion and data sources

The 2023 State of the Nation Report includes results for patients in England and Wales diagnosed with bowel cancer 1 April 2021 – 31 March 2022.

Adjuvant chemotherapy estimates include patients undergoing major resection for pathological stage III colon cancer between 01 December 2018 and 31 August 2021. Severe acute toxicity estimates include patients receiving adjuvant chemotherapy after major resection for pathological stage III colon cancer between 01 December 2018 and 31 August 2021.

APER/Hartmann's and unclosed ileostomy estimates include patients undergoing rectal cancer surgery between 01 April 2016 and 31 March 2021. Annual rectal cancer resection volume includes major resections for rectal cancer performed between 01 April 2021 and 31 March 2022.

30-day unplanned readmission, 30-day unplanned return to theatre and 90-day mortality are further restricted to patients undergoing surgery up to 31 October 2022. Two-year mortality estimates include patients undergoing a major resection between 1 April 2019 and 31 March 2020.

The Audit dataset is linked to Hospital Episode Statistics (HES) and Patient Episode Database for Wales (PEDW) at patient level to obtain further information on patient care and follow-up such as stoma reversal and emergency readmissions in England/Wales. Linkage to Office for National Statistics records provides information about date and cause of death.

NBOCA also links to the National Radiotherapy Dataset (RTDS) for information on radiotherapy treatment in England, and the Systemic Anti-Cancer Therapy database (SACT) for information on chemotherapy treatment in England.

Data Quality

Case ascertainment

Number of patients reported to the Audit as a percentage of the number of patients admitted for the first time to the trust/network with a diagnosis of bowel cancer within the audit period

according to HES in England and PEDW in Wales. This can be larger than 100 if more patients are reported to the Audit than identified in HES/PEDW.

Data completeness

% of relevant patient group with useable value of data item.

ASA grade

% of patients recorded as having a major resection who have a particular ASA grade recorded, or no ASA recorded.

Management of all patients

Clinical Nurse Specialist

% of patients recorded as having been seen by a clinical nurse specialist or a member of their team.

Management of patients having major resection

At least 12 lymph nodes excised (%)

% of colon cancer patients undergoing major resection with a recorded number of lymph nodes, who had at least 12 lymph nodes examined, reported by the trust/hospital/MDT providing major resection.

Adjuvant chemotherapy (%)

% of adjuvant chemotherapy in patients undergoing major resection for pathological stage III colon cancer between 01 June 2017 and 31 August 2020. These are unadjusted chemotherapy rates reported by the trust/hospital/MDT providing major resection. SACT and HES data are used for patients treated in England and PEDW data are used for patients treated in Wales.

Severe acute toxicity after adjuvant chemotherapy (%)

% risk-adjusted overnight admission for severe acute toxicity in patients receiving adjuvant chemotherapy for stage III colon cancer, reported by the trust/hospital/MDT providing chemotherapy. ICD-10 diagnosis codes in overnight admissions in HES/ PEDW from the first cycle of chemotherapy to 8 weeks after the last cycle of chemotherapy are used to identify severe acute toxicity. Estimates are risk-adjusted for age, sex, number of comorbidities, performance status and staging.

Rectal cancer patients

Neo-adjuvant therapy (%)

% of rectal cancer patients having short- or long-course radiotherapy prior to major resection, reported by the trust/hospital/MDT providing major resection.

Circumferential resection margin: Negative (%)

% of rectal cancer patients undergoing major resection whose CRM is reported to be negative, reported by the trust/hospital/MDT providing major resection.

Circumferential resection margin: Recorded (%)

% of rectal cancer patients undergoing a major resection who have a recorded CRM, reported by the trust/hospital/MDT providing major resection.

Rectal volume

Reported number of rectal major resections between 1 April 2021 and 31 March 2022, reported by the trust/hospital/MDT providing major resection.

APER/Hartmann's (%)

% of patients whose rectal cancer resection is an abdomino-perineal excision of rectum (APER)/pelvic exenteration/Hartmann's 1 April 2016 to 30 September 2021, reported by the trust/hospital/MDT providing major resection.

Compare trust outcomes

Trust outcomes are reported by the trust/hospital/MDT providing major resection. Funnel plots display trust risk-adjusted outcomes only for 2-year mortality this year. Other outcomes are not outlier-reported this year because they are measured on patients diagnosed and/or treated during the COVID-19 pandemic. The funnel regions represent the 95 per cent limit and the 99.8 per cent limit for trusts compared to the national average. Those trusts with results outside the outer (99.8 per cent) limit are considered potential outliers. Previous years of results show funnel plots for all five trust outcomes.

Risk adjustment is performed using the seven items listed under Data Quality as well as mode of admission (elective/emergency) and number of co-morbidities according to HES/PEDW, and an interaction between age and distant metastases. Missing values are imputed using Multiple Imputation. The model for two-year mortality additionally includes interactions between follow-up time (0-3 months after surgery vs. 3-24 months after surgery) and all of the risk factors.

See FAQs https://www.nboca.org.uk/about/fag/ for more details.

Data Quality

| All Patients: | Trust | Network | National |
|-----------------------------|-------|---------|----------|
| Number of patients in Audit | 368 | 1703 | 35779 |
| Case ascertainment (%) | Good | Good | Good |
| Data completeness of: | | | |
| – Pre-treatment TNM (%) | 89 | 86 | 86 |
| – Performance status (%) | 90 | 76 | 89 |

| Patients having major resection: | Trust | Network | National |
|---------------------------------------|-------|---------|----------|
| Number of patients in Audit | 161 | 821 | 19631 |
| ASA grade 1 (%) | 8 | 5 | 9 |
| ASA grade 2 (%) | 63 | 59 | 53 |
| ASA grade 3 (%) | 25 | 22 | 31 |
| ASA grade 4+ (%) | N/A | 1 | 2 |
| ASA grade not recorded (%) | 4 | 13 | 4 |
| Data completeness of: | | | |
| 7 Audit items for risk-adjustment (%) | 94 | 84 | 87 |

Key:> 80% 50 - 80% < 50%

Management of all patients

| All Patients: | Trust | Network | National |
|---------------------------------------|-------|---------|----------|
| Number of patients in Audit | 368 | 1703 | 35779 |
| Seen by Clinical Nurse Specialist (%) | 43 | 76 | 88 |

Management of patients having major resection

| Patients having major resection: | Trust | Network | National |
|-------------------------------------|-------|---------|----------|
| Number of patients in Audit | 161 | 821 | 19614 |
| Distant metastases (%) | 4 | 5 | 5 |
| Urgent or emergency surgery (%) | 6 | 20 | 15 |
| At least 12 lymph nodes excised (%) | 92 | 90 | 84 |

| Patients having major resection: | Trust | Network | National |
|---|-------|---------|----------|
| Laparoscopic surgery attempted (%) | 97 | 81 | 72 |
| Risk-adjusted length of stay > 5 days (%) | 53 | 63 | 55 |

| Patients having major resection for stage III colon cancer: | Trust | Network | National |
|---|-------|---------|----------|
| Number of patients in audit | 84 | N/A | 10747 |
| Adjuvant chemotherapy (%) | 64 | N/A | 62 |


| Patients receiving chemotherapy after major resection for stage III colon cancer: | Trust | Network | National |
|---|-------|---------|----------|
| Number of patients in audit | 55 | N/A | 6608 |
| Severe acute toxicity after adjuvant chemotherapy (%) | 19 | N/A | 22 |

Rectal cancer patients

| Patients having major resection: | Trust | Network | National |
|----------------------------------|-------|---------|----------|
| Number of patients in Audit | 27 | N/A | 4251 |
| Neoadjuvant therapy (%) | 54 | N/A | 33 |

| Patients having major resection: | Trust | Network | National |
|--|-------|---------|----------|
| Circumferential resection margin: Recorded (%) | 93 | 96 | 89 |
| Circumferential resection margin: Negative (%) | 93 | 90 | 83 |
| Rectal volume | 30 | N/A | 4377 |

| Patients in APER/Hartmanns estimate: | Trust | Network | National |
|---|-------|---------|----------|
| Number of patients in APER/Hartmanns estimate | 118 | 909 | 19644 |
| APER/Hartmanns (%) | 36 | 31 | 37 |



| Trust | Number | Adjusted | Observed | |
|--|--------|----------|----------|--|
| Calderdale and Huddersfield NHS Foundation Trust | 118 | 19.2% | 15.8% | |
| Other trusts within the region: West Yorkshire and Harrogate | | | | |
| Airedale NHS Foundation Trust | 78 | 14.5% | 15.4% | |
| Bradford Teaching Hospitals NHS Foundation Trust | 119 | 20% | 15.7% | |

| Trust | Number | Adjusted | Observed |
|---|--------|----------|----------|
| Harrogate and District NHS Foundation Trust | 81 | 11.1% | 10.6% |
| Leeds Teaching Hospitals NHS Trust | 267 | 11.1% | 10.5% |
| Mid Yorkshire Hospitals NHS Trust | 198 | 13.2% | 13.1% |