

University Hospital Southampton NHS Foundation Trust

Wessex A16

The University Hospital Southampton NHS Foundation Trust incorporates Southampton General Hospital <http://www.uhs.nhs.uk> <<http://www.uhs.nhs.uk>>

Patient inclusion and data sources

The majority of these results are for patients in England and Wales diagnosed with bowel cancer 1 Apr 2017 – 31 Mar 2018. The exceptions are for adjuvant chemotherapy and the four trust outcomes: 30-day unplanned readmission, 90-day mortality, 2-year mortality and 18-month stoma rate. 30-day unplanned readmission and 90-day mortality are further restricted to patients undergoing surgery up to 31 October 2018 for patients treated in England, and up to 28 February 2018 for patients treated in Wales. Two-year mortality estimates include patients undergoing a major resection between 1 April 2015 and 31 March 2016. 18-month stoma rates include rectal cancer patients undergoing a major resection between 1 April 2014 and 31 March 2017. Adjuvant chemotherapy estimates include patients undergoing major resection for pathological stage III colon cancer between 01 June 2014 and 31 August 2017 in England only.

The Audit dataset is linked to Hospital Episode Statistics (HES) and Patient Episode Database for Wales (PEDW) at the patient level to obtain further information on patient care and follow-up such as stoma reversal and emergency readmissions in England/Wales. The dataset also links to Office of National Statistics records. These provide information about date, place and cause of death.

NBOCA also links to the National Radiotherapy Dataset (RTDS) for information on radiotherapy treatment in England, and the Systemic Anti-Cancer Therapy database (SACT) for information on chemotherapy treatment in England.

Data Quality

Case ascertainment

Number of patients reported to the Audit as a percentage of the number of patients admitted for the first time to the trust/network with a diagnosis of bowel cancer within the audit period according to HES. This can be larger than 100 if more patients are reported to the Audit than identified in HES.

Data completeness

% of relevant patient group with useable value of data item.

ASA grade

% of patients recorded as having a major resection who have a particular ASA grade recorded, or no ASA recorded.

Management of all patients

Potentially curative patients

Patients electively diagnosed with colon cancer with pre-treatment staging of T2-T4 and no evidence of metastatic disease.

Management of patients having major resection

At least 12 lymph nodes excised (%)

% of colon cancer patients undergoing major resection with a recorded number of lymph nodes, who had at least 12 lymph nodes examined.

Adjuvant chemotherapy

% of adjuvant chemotherapy in patients undergoing major resection for pathological stage III colon cancer between 01 June 2014 and 31 August 2017 in England only. These are unadjusted chemotherapy rates.

Rectal cancer patients

Neo-adjuvant therapy (%)

% of rectal cancer patients having short- or long-course radiotherapy prior to major resection.

Circumferential resection margin: Negative (%)

% of rectal cancer patients undergoing major resection whose CRM is reported to be negative.

Circumferential resection margin: Recorded (%)

% of rectal cancer patients undergoing a major resection who have a recorded CRM.

APER rate (%)

% of patients with rectal cancer undergoing abdominoperineal excision of the rectum and therefore having a permanent stoma.

Compare trust outcomes

Funnel plots display trust risk-adjusted outcomes for 90-day mortality, 30-day unplanned readmission, 2-year mortality and, for rectal cancer patients, 18-month stoma rate. The funnel regions represent the 95 per cent limit and the 99.8 per cent limit for trusts compared to the national average. Those trusts with results outside the outer (99.8 per cent) limit are considered potential outliers.

Risk adjustment is performed using the seven items listed under Data Quality as well as mode of admission (elective/emergency) and number of co-morbidities according to HES/PEDW, and an interaction between age and distant metastases. Missing values are imputed using Multiple Imputation. The model for two-year mortality additionally includes interactions between follow-up time (0-3 months after surgery vs. 3-24 months after surgery) and all of the risk factors.

A stoma is considered to be reversed if a HES/PEDW record with relevant code is identified within 18 months of the initial surgical procedure.

See [FAQs <https://www.nboca.org.uk/about/faq/>](https://www.nboca.org.uk/about/faq/) for more details.




Data Quality

All Patients:	Trust	Network	National
Number of patients in Audit	250	1475	31676
Case ascertainment (%)	85	96	95
Data completeness of:			

All Patients:	Trust	Network	National
– Pre-treatment TNM (%)	100	88	80
– Performance status (%)	100	76	86

Patients having major resection:	Trust	Network	National
Number of patients in Audit	176	893	19218
ASA grade 1 (%)	7	8	11
ASA grade 2 (%)	65	64	53
ASA grade 3 (%)	23	24	28
ASA grade 4+ (%)	5	2	3
ASA grade not recorded (%)	1	2	5
Data completeness of:			
7 Audit items for risk-adjustment (%)	99	94	86

Key:

	> 80%
	50 - 80%
	< 50%

Management of all patients

All Patients:	Trust	Network	National
Number of patients in Audit	250	1475	31676
Seen by Clinical Nurse Specialist (%)	93	95	87

All patients deemed potentially curative	Trust	Network	National
Number of potentially curative patients	70	412	7416
Undergoing major resection (%)	86	85	86

Management of patients having major resection


Patients having major resection:	Trust	Network	National
Number of patients in Audit	176	893	19218
Distant metastases (%)	6	6	8
Urgent or emergency surgery (%)	13	11	15
At least 12 lymph nodes excised (%)	89	88	84
Laparoscopic surgery attempted (%)	72	82	69
Risk-adjusted length of stay > 5 days (%)	63	57	62

Patients having major resection for stage III colon cancer:	Trust	Network	National
Number of patients in audit	124	614	11837
Adjuvant chemotherapy (%)	52	57	62


Rectal cancer patients

Patients having major resection:	Trust	Network	National
Number of patients in Audit	51	246	4588
Neoadjuvant therapy (%)	46	22	33
Circumferential resection margin: Recorded (%)	98	96	90
Circumferential resection margin: Negative (%)	94	84	80
Rectal volume	N/A	N/A	N/A


Patients in APER/Hartmanns estimate:	Trust	Network	National
Number of patients in APER/Hartmanns estimate	N/A	N/A	N/A
APER/Hartmanns (%)	N/A	N/A	N/A




Trust	Number	Adjusted	Observed
University Hospital Southampton NHS Foundation Trust	159	8.8%	8.8%
Other trusts within the region: Wessex			
Dorset County Hospital NHS Foundation Trust	82	12.6%	12.2%
Hampshire Hospitals NHS Foundation Trust - Basingstoke and North Hampshire Hospital	74	21.1%	21.6%
Hampshire Hospitals NHS Foundation Trust - Royal Hampshire County Hospital	78	5.4%	5.1%
Isle of Wight NHS Trust	55	12.8%	12.7%
Poole Hospital NHS Foundation Trust	91	14.5%	14.3%
Portsmouth Hospitals NHS Trust	182	13%	13.2%
The Royal Bournemouth and Christchurch Hospitals NHS Foundation Trust	89	5.7%	5.6%



Trust	Number	Adjusted	Observed
University Hospital Southampton NHS Foundation Trust	173	3.7%	3.5%
Other trusts within the region: Wessex			
Dorset County Hospital NHS Foundation Trust	85	1.7%	2.4%
Hampshire Hospitals NHS Foundation Trust - Basingstoke and North Hampshire Hospital	82	2.1%	1.2%
Hampshire Hospitals NHS Foundation Trust - Royal Hampshire County Hospital	79	1.5%	1.3%
Isle of Wight NHS Trust	56	4.5%	3.6%
Poole Hospital NHS Foundation Trust	98	2.7%	3.1%
Portsmouth Hospitals NHS Trust	190	0.8%	0.5%
The Royal Bournemouth and Christchurch Hospitals NHS Foundation Trust	94	1%	1.1%



Trust	Number	Adjusted	Observed
University Hospital Southampton NHS Foundation Trust	171	18.7%	13.7%
Other trusts within the region: Wessex			
Dorset County Hospital NHS Foundation Trust	90	15.1%	15.6%
Hampshire Hospitals NHS Foundation Trust - Basingstoke and North Hampshire Hospital	73	11.6%	8.6%
Hampshire Hospitals NHS Foundation Trust - Royal Hampshire County Hospital	94	27.2%	25.4%
Isle of Wight NHS Trust	69	22.3%	21.5%
Poole Hospital NHS Foundation Trust	94	7.7%	8.9%
Portsmouth Hospitals NHS Trust	201	15.6%	16.8%
The Royal Bournemouth and Christchurch Hospitals NHS Foundation Trust	121	14.7%	12.4%



Trust	Number	Adjusted	Observed
University Hospital Southampton NHS Foundation Trust	148	37.3%	37.2%
Other trusts within the region: Wessex			
Dorset County Hospital NHS Foundation Trust	57	45.6%	45.6%
Hampshire Hospitals NHS Foundation Trust - Basingstoke and North Hampshire Hospital	57	34.9%	35.1%
Hampshire Hospitals NHS Foundation Trust - Royal Hampshire County Hospital	39	42.9%	46.2%
Isle of Wight NHS Trust	47	46.6%	44.7%
Poole Hospital NHS Foundation Trust	70	39%	40%
Portsmouth Hospitals NHS Trust	196	43.6%	43.4%
The Royal Bournemouth and Christchurch Hospitals NHS Foundation Trust	91	51.1%	50.6%