

#### Imperial College Healthcare NHS Trust

**RM** Partners A17

The Imperial College Healthcare NHS Trust incorporates Charing Cross Hospital, St Mary's Hospital Paddington, and Hammersmith Hospital in London <a href="https://www.imperial.nhs.uk/">https://www.imperial.nhs.uk/</a> <a href="https://www.imperial

#### Patient inclusion and data sources

The 2023 State of the Nation Report includes results for patients in England and Wales diagnosed with bowel cancer 1 April 2021 – 31 March 2022.

Adjuvant chemotherapy estimates include patients undergoing major resection for pathological stage III colon cancer between 01 December 2018 and 31 August 2021. Severe acute toxicity estimates include patients receiving adjuvant chemotherapy after major resection for pathological stage III colon cancer between 01 December 2018 and 31 August 2021.

APER/Hartmann's and unclosed ileostomy estimates include patients undergoing rectal cancer surgery between 01 April 2016 and 31 March 2021. Annual rectal cancer resection volume includes major resections for rectal cancer performed between 01 April 2021 and 31 March 2022.

30-day unplanned readmission, 30-day unplanned return to theatre and 90-day mortality are further restricted to patients undergoing surgery up to 31 October 2022. Two-year mortality estimates include patients undergoing a major resection between 1 April 2019 and 31 March 2020.

The Audit dataset is linked to Hospital Episode Statistics (HES) and Patient Episode Database for Wales (PEDW) at patient level to obtain further information on patient care and follow-up such as stoma reversal and emergency readmissions in England/Wales. Linkage to Office for National Statistics records provides information about date and cause of death.

NBOCA also links to the National Radiotherapy Dataset (RTDS) for information on radiotherapy treatment in England, and the Systemic Anti-Cancer Therapy database (SACT) for information on chemotherapy treatment in England.

#### **Data Quality**

#### Case ascertainment

Number of patients reported to the Audit as a percentage of the number of patients admitted for the first time to the trust/network with a diagnosis of bowel cancer within the audit period

according to HES in England and PEDW in Wales. This can be larger than 100 if more patients are reported to the Audit than identified in HES/PEDW.

#### **Data completeness**

% of relevant patient group with useable value of data item.

#### **ASA** grade

% of patients recorded as having a major resection who have a particular ASA grade recorded, or no ASA recorded.

#### Management of all patients

#### **Clinical Nurse Specialist**

% of patients recorded as having been seen by a clinical nurse specialist or a member of their team.

#### Management of patients having major resection

#### At least 12 lymph nodes excised (%)

% of colon cancer patients undergoing major resection with a recorded number of lymph nodes, who had at least 12 lymph nodes examined, reported by the trust/hospital/MDT providing major resection.

#### Adjuvant chemotherapy (%)

% of adjuvant chemotherapy in patients undergoing major resection for pathological stage III colon cancer between 01 June 2017 and 31 August 2020. These are unadjusted chemotherapy rates reported by the trust/hospital/MDT providing major resection. SACT and HES data are used for patients treated in England and PEDW data are used for patients treated in Wales.

#### Severe acute toxicity after adjuvant chemotherapy (%)

% risk-adjusted overnight admission for severe acute toxicity in patients receiving adjuvant chemotherapy for stage III colon cancer, reported by the trust/hospital/MDT providing chemotherapy. ICD-10 diagnosis codes in overnight admissions in HES/ PEDW from the first cycle of chemotherapy to 8 weeks after the last cycle of chemotherapy are used to identify severe acute toxicity. Estimates are risk-adjusted for age, sex, number of comorbidities, performance status and staging.

#### **Rectal cancer patients**

#### **Neo-adjuvant therapy (%)**

% of rectal cancer patients having short- or long-course radiotherapy prior to major resection, reported by the trust/hospital/MDT providing major resection.

#### **Circumferential resection margin: Negative (%)**

% of rectal cancer patients undergoing major resection whose CRM is reported to be negative, reported by the trust/hospital/MDT providing major resection.

#### **Circumferential resection margin: Recorded (%)**

% of rectal cancer patients undergoing a major resection who have a recorded CRM, reported by the trust/hospital/MDT providing major resection.

#### **Rectal volume**

Reported number of rectal major resections between 1 April 2021 and 31 March 2022, reported by the trust/hospital/MDT providing major resection.

#### APER/Hartmann's (%)

% of patients whose rectal cancer resection is an abdomino-perineal excision of rectum (APER)/pelvic exenteration/Hartmann's 1 April 2016 to 30 September 2021, reported by the trust/hospital/MDT providing major resection.

#### **Compare trust outcomes**

Trust outcomes are reported by the trust/hospital/MDT providing major resection. Funnel plots display trust risk-adjusted outcomes only for 2-year mortality this year. Other outcomes are not outlier-reported this year because they are measured on patients diagnosed and/or treated during the COVID-19 pandemic. The funnel regions represent the 95 per cent limit and the 99.8 per cent limit for trusts compared to the national average. Those trusts with results outside the outer (99.8 per cent) limit are considered potential outliers. Previous years of results show funnel plots for all five trust outcomes.

Risk adjustment is performed using the seven items listed under Data Quality as well as mode of admission (elective/emergency) and number of co-morbidities according to HES/PEDW, and an interaction between age and distant metastases. Missing values are imputed using Multiple Imputation. The model for two-year mortality additionally includes interactions between follow-up time (0-3 months after surgery vs. 3-24 months after surgery) and all of the risk factors.

See FAQs <a href="https://www.nboca.org.uk/about/fag/">https://www.nboca.org.uk/about/fag/</a> for more details.

### **Data Quality**

All Patients:	Trust	Network	National
Number of patients in Audit	231	1558	35779
Case ascertainment (%)	Good	Fair	Good
Data completeness of:			
– Pre-treatment TNM (%)	89	81	86
– Performance status (%)	81	78	89

Patients having major resection:	Trust	Network	National
Number of patients in Audit	128	848	19631
ASA grade 1 (%)	91	22	9
ASA grade 2 (%)	6	52	53
ASA grade 3 (%)	1	21	31
ASA grade 4+ (%)	N/A	1	2
ASA grade not recorded (%)	2	4	4
Data completeness of:			
7 Audit items for risk-adjustment (%)	91	93	87

# **Key:**> 80% 50 - 80% < 50%

## Management of all patients

All Patients:	Trust	Network	National
Number of patients in Audit	231	1558	35779
Seen by Clinical Nurse Specialist (%)	91	85	88

## Management of patients having major resection

Patients having major resection:	Trust	Network	National
Number of patients in Audit	128	848	19614
Distant metastases (%)	3	6	5
Urgent or emergency surgery (%)	12	10	15
At least 12 lymph nodes excised (%)	92	93	84

Patients having major resection:	Trust	Network	National
Laparoscopic surgery attempted (%)	53	63	72
Risk-adjusted length of stay > 5 days (%)	84	74	55

Patients having major resection for stage III colon cancer:	Trust	Network	National
Number of patients in audit	66	N/A	10747
Adjuvant chemotherapy (%)	65	N/A	62


Patients receiving chemotherapy after major resection for stage III colon cancer:	Trust	Network	National
Number of patients in audit	70	N/A	6608
Severe acute toxicity after adjuvant chemotherapy (%)	22	N/A	22

## Rectal cancer patients

Patients having major resection:	Trust	Network	National
Number of patients in Audit	21	N/A	4251
Neoadjuvant therapy (%)	20	N/A	33

Patients having major resection:	Trust	Network	National
Circumferential resection margin: Recorded (%)	95	98	89
Circumferential resection margin: Negative (%)	90	94	83
Rectal volume	24	N/A	4377

Patients in APER/Hartmanns estimate:	Trust	Network	National
Number of patients in APER/Hartmanns estimate	111	985	19644
APER/Hartmanns (%)	27	31	37



Trust	Number	Adjusted	Observed	
Imperial College Healthcare NHS Trust	131	18.8%	16.4%	
Other trusts within the region: RM Partners				
Chelsea and Westminster Hospital NHS Foundation Trust	127	13.7%	15.9%	
Croydon Health Services NHS Trust	56	13.3%	9.5%	

Trust	Number	Adjusted	Observed
Epsom and St Helier University Hospitals NHS Trust	123	15.9%	13.2%
Kingston Hospital NHS Foundation Trust	102	14.8%	17.5%
London North West Hospitals NHS Trust	238	18.3%	17%
The Hillingdon Hospitals NHS Foundation Trust	71	15.5%	21.8%
The Royal Marsden NHS Foundation Trust	48	10.3%	8.7%