

Northumbria Healthcare NHS Foundation Trust

Northern A09

The Northumbria Healthcare NHS Foundation Trust incorporates Hexham General Hospital, Wansbeck Hospital and North Tyneside General Hospital <u>https://www.northumbria.nhs.uk/</u> <https://www.northumbria.nhs.uk/>

Patient inclusion and data sources

The majority of these results are for patients in England and Wales diagnosed with bowel cancer 1 Apr 2018 – 31 Mar 2019. The exceptions are for adjuvant chemotherapy and the four trust outcomes: 30-day unplanned readmission, 90-day mortality and 2-year mortality. 30-day unplanned readmission and 90-day mortality are further restricted to patients undergoing surgery up to 31 October 2019. Two-year mortality estimates include patients undergoing a major resection between 1 April 2016 and 31 March 2017. Adjuvant chemotherapy estimates include patients undergoing major resection for pathological stage III colon cancer between 01 December 2015 and 31 August 2018.

The Audit dataset is linked to Hospital Episode Statistics (HES) and Patient Episode Database for Wales (PEDW) at the patient level to obtain further information on patient care and follow-up such as stoma reversal and emergency readmissions in England/Wales. The dataset also links to Office of National Statistics records. These provide information about date, place and cause of death.

NBOCA also links to the National Radiotherapy Dataset (RTDS) for information on radiotherapy treatment in England, and the Systemic Anti-Cancer Therapy database (SACT) for information on chemotherapy treatment in England.

Data Quality

Case ascertainment

Number of patients reported to the Audit as a percentage of the number of patients admitted for the first time to the trust/network with a diagnosis of bowel cancer within the audit period according to NCRAS in England and PEDW in Wales. This can be larger than 100 if more patients are reported to the Audit than identified in NCRAS/PEDW.

Data completeness

% of relevant patient group with useable value of data item.

ASA grade

% of patients recorded as having a major resection who have a particular ASA grade recorded, or no ASA recorded.

Management of all patients

Potentially curative patients

Patients electively diagnosed with colon cancer with pre-treatment staging of T2-T4 and no evidence of metastatic disease.

Management of patients having major resection

At least 12 lymph nodes excised (%)

% of colon cancer patients undergoing major resection with a recorded number of lymph nodes, who had at least 12 lymph nodes examined.

Adjuvant chemotherapy

% of adjuvant chemotherapy in patients undergoing major resection for pathological stage III colon cancer between 01 December 2015 and 31 August 2018 in England only. These are unadjusted chemotherapy rates.

Rectal cancer patients

Neo-adjuvant therapy (%)

% of rectal cancer patients having short- or long-course radiotherapy prior to major resection.

Circumferential resection margin: Negative (%)

% of rectal cancer patients undergoing major resection whose CRM is reported to be negative.

Circumferential resection margin: Recorded (%)

% of rectal cancer patients undergoing a major resection who have a recorded CRM.

APER rate (%)

% of patients with rectal cancer undergoing abdominoperineal excision of the rectum and therefore having a permanent stoma.

Compare trust outcomes

Funnel plots display trust risk-adjusted outcomes for 90-day mortality, 30-day unplanned readmission and 2-year mortality. This year we are not reporting 18-month stoma rates in readiness for two new stoma indicators in 2021. The funnel regions represent the 95 per cent limit and the 99.8 per cent limit for trusts compared to the national average. Those trusts with results outside the outer (99.8 per cent) limit are considered potential outliers.

Risk adjustment is performed using the seven items listed under Data Quality as well as mode of admission (elective/emergency) and number of co-morbidities according to HES/PEDW, and an interaction between age and distant metastases. Missing values are imputed using Multiple Imputation. The model for two-year mortality additionally includes interactions between follow-up time (0-3 months after surgery vs. 3-24 months after surgery) and all of the risk factors.

See FAQs <<u>https://www.nboca.org.uk/about/faq/></u> for more details.

Data Quality

All Patients:	Trust	Network	National
Number of patients in Audit	316	2011	29766
Case ascertainment (%)	100	99	90
Data completeness of:			
– Pre-treatment TNM (%)	82	75	83
– Performance status (%)	89	77	85

Patients having major resection:	Trust	Network	National
Number of patients in Audit	171	1128	16499
ASA grade 1 (%)	9	8	10
ASA grade 2 (%)	58	48	53
ASA grade 3 (%)	30	33	30
ASA grade 4+ (%)	3	4	3
ASA grade not recorded (%)	0	8	4
Data completeness of:			
7 Audit items for risk-adjustment (%)	100	77	86

Key:



Management of all patients

All Patients:	Trust	Network	National
Number of patients in Audit	316	2011	29766
Seen by Clinical Nurse Specialist (%)	80	82	86

All patients deemed potentially curative	Trust	Network	National
Number of potentially curative patients	97	523	7226
Undergoing major resection (%)	76	85	86

Management of patients having major resection

Patients having major resection:	Trust	Network	National
Number of patients in Audit	171	1128	16474


Patients having major resection:	Trust	Network	National
Distant metastases (%)	7	5	6
Urgent or emergency surgery (%)	12	15	15
At least 12 lymph nodes excised (%)	84	74	84
Laparoscopic surgery attempted (%)	72	85	72
Risk-adjusted length of stay > 5 days (%)	43	52	62

Patients having major resection for stage III colon cancer:	Trust	Network	National
Number of patients in audit	97	692	10641
Adjuvant chemotherapy (%)	60	58	61


Rectal cancer patients

Patients having major resection:	Trust	Network	National
Number of patients in Audit	44	279	3899
Neoadjuvant therapy (%)	24	32	32
Circumferential resection margin: Recorded (%)	100	73	86
Circumferential resection margin: Negative (%)	95	70	79
Rectal volume	N/A	N/A	N/A


Patients in APER/Hartmanns estimate:	Trust	Network	National
Number of patients in APER/Hartmanns estimate	N/A	N/A	N/A
APER/Hartmanns (%)	N/A	N/A	N/A



Trust	Number	Adjusted	Observed
Northumbria Healthcare NHS Foundation Trust	168	13.2%	13.1%
Other trusts within t	he region: Northern		
County Durham and Darlington NHS Foundation Trust	196	10.5%	10.7%
Gateshead Health NHS Foundation Trust	121	11%	10.7%
North Tees and Hartlepool NHS Foundation Trust	122	9.9%	9.8%
South Tyneside and Sunderland NHS Foundation Trust - South Tyneside District Hospital	54	15.1%	14.8%
South Tyneside and Sunderland NHS Foundation Trust - Sunderland Royal Hospital	118	16.6%	17%
The Newcastle Upon Tyne Hospitals NHS Foundation Trust	146	8.6%	8.9%



Trust	Number	Adjusted	Observed
Northumbria Healthcare NHS Foundation Trust	171	3.1%	2.9%
Other trusts within t	he region: Northern	L	
County Durham and Darlington NHS Foundation Trust	204	1.5%	1.5%
Gateshead Health NHS Foundation Trust	122	2.2%	2.5%
North Tees and Hartlepool NHS Foundation Trust	125	3.7%	5.6%
South Tyneside and Sunderland NHS Foundation Trust - South Tyneside District Hospital	58	1.9%	1.7%
South Tyneside and Sunderland NHS Foundation Trust - Sunderland Royal Hospital	120	2.6%	2.5%
The Newcastle Upon Tyne Hospitals NHS Foundation Trust	151	3.4%	3.3%



Trust	Number	Adjusted	Observed
Northumbria Healthcare NHS Foundation Trust	155	14.2%	14.6%
Other trusts within	the region: North	iern	
County Durham and Darlington NHS Foundation Trust	170	14.7%	14.4%
Gateshead Health NHS Foundation Trust	90	14.2%	16.7%
North Cumbria University Hospitals NHS Trust	75	18.3%	17.6%
North Tees and Hartlepool NHS Foundation Trust	142	17.9%	20.9%
South Tees Hospitals NHS Foundation Trust	199	21.6%	18.6%
South Tyneside and Sunderland NHS Foundation Trust - South Tyneside District Hospital	55	11.1%	18.3%
South Tyneside and Sunderland NHS Foundation Trust - Sunderland Royal Hospital	105	21.9%	21.7%
The Newcastle Upon Tyne Hospitals NHS Foundation Trust	166	19.2%	17.8%